

## **POST-OPERATIVE INSTRUCTIONS FOR CHIN AUGMENTATION**

1. The incision is either located on the inside aspect of the lower lip or in the skin underneath the chin. For incisions located on the skin, it is usually covered with tape. However, if the tape falls off, then clean the suture line 4 times daily with hydrogen peroxide on a Q-tip, gently rolling the Q-tip over the suture line to remove any dried blood, then apply a thin layer of Vaseline or Bacitracin antibiotic ointment. It is not necessary to continue these cleanings throughout the night. For incisions located within the inside aspect of the lower lip, gently brush your teeth with a baby toothbrush and rinse with Peridex mouthwash 15 mL three times a day and after meals. Avoid excessive stretching of the lower lip.
2. Clean all the suture lines 4 times daily with hydrogen peroxide on a Q-tip, gently rolling the Q-tip over the suture line to remove any dried blood, then apply a thin layer of Vaseline or Bacitracin antibiotic ointment.
3. It is common for the swelling and bruising to increase in the neck region 2-3 days following surgery. It is strongly recommended that you use cold compresses (wash cloths soaked in ice water) during the first 24-48 hours to help keep the swelling to a minimum. Please do not directly apply ice to the area.
4. Watch for any abnormal swelling in the neck region shaped like a “goose egg” that feels firm or fluid filled. If this happens, please contact us at 949.650.8882, as this may indicate a collection of fluid under the skin and may need to be drained by Dr. Batniji.
5. The chin area may have tape in place. Dr. Batniji will remove the tape one week after surgery. However, you are able to shower as you normally would on the first day following surgery.
6. Keep your head elevated at all times for the first week of recovery. We recommend sleeping on 2-3 pillows at night.
7. Avoid turning your head from side to side. Turn with your whole upper body.
8. Please do not bend forward, strain or lift anything during the first week of recovery.

9. Shower at least once a day, shampooing your hair (with baby shampoo only). Prior to showering, remove the wrap dressing that Dr. Batniji placed in the office; this is an excellent opportunity to evaluate the neck for any fluid collections. After showering, replace the wrap dressing; Dr. Batniji and staff will show you how to apply the dressing on your office visit the day after your procedure. If the dressing loosens during the day, please replace it. It is important that the dressing is smooth against the skin and snug to minimize the potential for contour irregularities and/or fluid collection formations.
10. **DO NOT TAKE ASPIRIN or IBUPROFEN PRODUCTS AS THESE MAY INCREASE THE RISK OF BLEEDING.**
11. Do not take narcotic medication on an empty stomach as this may make you feel nauseous.
12. It is helpful to wean yourself off of the pain medication as soon as possible. Try taking Tylenol in place of the narcotic medication. Please be aware of the maximum recommended daily dose of acetaminophen (active ingredient in both Tylenol and narcotic pain medication).
13. Dr. Batniji will place a neck wrap on you immediately following your procedure. The day after your procedure, Dr. Batniji will remove this neck wrap and replace it with a lighter wrap. You are to wear this wrap at all times for 1 week after the procedure. You may remove this wrap to shower. After the first week, Dr. Batniji may instruct you to wear the wrap dressing at night for another 2-3 weeks.
14. Dr. Batniji will remove sutures and staples one week after surgery.

If you have any further questions or concerns, please do not hesitate to contact our office at 949.650.8882.