

361 HOSPITAL RD., #329 · NEWPORT BEACH, CA 92663 · 949.650.8882 / 949.650.2293 FAX  
416 N. BEDFORD DR., #200 · BEVERLY HILLS, CA 90210 · 310.467.2180 / 949.650.2293 FAX  
WWW.DRBATNIJI.COM · @RAMIKBATNIJI · INFO@DRBATNIJI.COM

*Your recovery is just as important as your surgery.*

## POST-OPERATIVE INSTRUCTIONS

### *Browlift Surgery*

Following these instructions closely will help you heal comfortably and achieve the best possible result. Please read the entire document and keep it handy throughout your recovery. Call our office at **949.650.8882** (Newport Beach) or **310.467.2180** (Beverly Hills) with any questions or concerns.

### CALL OUR OFFICE IMMEDIATELY

**Newport Beach:** 949.650.8882    **Beverly Hills:** 310.467.2180

Contact us right away if you experience any of the following:

- New or increasing swelling on the forehead, brow, or scalp
- A firm or fluid-filled area developing under the skin
- Active bleeding from an incision that is controlled with light pressure but recurs
- Shortness of breath, or calf pain/swelling

**For life-threatening symptoms,** always call 911 or go to the nearest emergency room first.

### What to Expect During Recovery

The following sensations are common and expected during early healing, and typically improve over days to a few weeks:

- Headache
- Numbness of the forehead and scalp
- Mild asymmetry during early healing (the two sides often swell at slightly different rates)
- Temporary contour irregularities that smooth out over weeks to a few months
- Tingling, itching, and occasional brief “electric zinger” sensations as sensation nerves heal

## Wound Care

---

- Open a fresh bottle of hypochlorous acid around the time of surgery for best potency.
- **Showers at least once a day**, shampooing your hair with a gentle shampoo (e.g., baby shampoo). After showering, spray the incision lines directly with hypochlorous acid spray and allow it to dry for 1 minute.
- Do not pick at crusts along the incision line. Allow the crusts to soften with the shower and the hypochlorous acid treatments.
- Do not submerge incisions (no baths, pools, hot tubs, or ocean) for 3–4 weeks.

## Swelling, Bruising, and Head Positioning

---

- Swelling and bruising often peak 2–3 days after surgery. This is expected.
- Apply cold compresses (for example, a washcloth soaked in ice water) over the forehead and brow. Do not apply ice directly to the skin.
- Keep your head elevated at all times during the first 2 weeks. Sleep on 2–3 pillows or in a recliner. A travel neck pillow may help maintain a neutral head position during sleep.
- Sleep on your back, not your side, for approximately 2–4 weeks after surgery. Side sleeping can distort healing and cause asymmetry.
- Temporary fullness of the forehead may represent normal post-operative swelling or a small fluid collection. If it is a small fluid collection, it can be easily managed in the office if needed. Please call us if you feel a distinct firm or fluid-filled area.
- Most visible swelling improves over 2–3 weeks, but subtle swelling can persist for a few months as the tissues continue to settle.

## Blood Pressure, Nausea, and Vomiting

---

Keeping your blood pressure well controlled in the first 2 weeks is one of the most important things you can do to minimize a hematoma or fluid collection. Vomiting and forceful retching can significantly raise pressure in the face and neck and increase the risk of bleeding.

- Take all prescribed blood pressure medications exactly as directed.
- Avoid stress, straining, and sudden exertion.
- Take anti-nausea medication as prescribed, even preventively if you feel queasy.
- Take pain medications with food or a small snack to minimize nausea that can occur when taking pain medication on an empty stomach.

## Activity and Recovery Timeline

---

- Take short, gentle walks around the house starting the day of surgery. This helps reduce the risk of blood clots.
- For the first 2 weeks, avoid bending forward, lifting anything heavier than 5 pounds, and any form of straining — including heavy coughing, forceful sneezing, and excessive straining with bowel movements.
- Avoid sexual activity for 2 weeks, as this can raise blood pressure and increase the risk of bleeding.

- **Weeks 1–2:** you may gradually resume light, non-strenuous activity, as tolerated. **Avoid any activity that increases your heart rate or blood pressure, as that may increase the risk of bruising, swelling, and/or bleeding.**
- **Week 3 onward:** you may gradually resume strenuous activity. Begin slowly and work up to more strenuous activity. It is normal to experience some temporary swelling and tightness with increased activity at this stage.
- Do not drive while taking narcotic pain medication. You may drive once you are off narcotic pain medication and can comfortably turn your upper body to check blind spots.

## Medications and Pain Control

---

- **DO NOT TAKE ASPIRIN, IBUPROFEN, OR OTHER NSAIDs** (such as Advil, Motrin, Aleve, Naproxen, or Excedrin) during the 2 weeks before surgery or the 2 weeks after surgery. These can increase bleeding risk. If you are on Aspirin under the guidance of a cardiologist, you must ask your cardiologist if it is safe to be off Aspirin for your elective cosmetic procedure.
- Tylenol (Acetaminophen) is safe and a good medication for discomfort after surgery. Do not exceed 3,000 mg of Acetaminophen in 24 hours and remember that most narcotic pain medications already contain Acetaminophen — do not double-dose.
- We may prescribe Celebrex (Celecoxib) to take twice a day for the first week after surgery. Take this on a fixed schedule, not as needed, even on days you feel comfortable. Celebrex is a COX-2 selective anti-inflammatory that reduces pain and swelling without the bleeding risk of ibuprofen or aspirin. Taking it consistently helps you need less narcotic pain medication.
- If you experience discomfort, take Ativan (Lorazepam) as prescribed. This may help alleviate your discomfort.
- If you are still experiencing pain after taking Celebrex, Tylenol, and Ativan, you may take the prescribed narcotic pain medication. Take it with food to minimize nausea and wean off the narcotic pain medication as soon as possible.
- To minimize constipation from pain medication, stay hydrated, eat fiber-rich foods, and use a stool softener (such as Colace or Docusate) as needed while taking narcotic pain medication.
- Take any antibiotics prescribed by Dr. Batniji until the full course is finished.

## Diet

---

- Start with light, soft foods the day of surgery. Soft foods (yogurt, eggs, pasta, soups, smoothies) are best for the first day. We recommend a diet that is high in protein to help with the healing process.
- Advance to your normal diet as tolerated.
- Stay well hydrated.
- Avoid very salty foods for the first few weeks after surgery, as excess sodium can worsen swelling.

- Include fiber-rich foods to help minimize constipation.
- We recommend using throat lozenges such as Cepacol or Ricola if you have a sore throat.

## Nicotine, Alcohol, and Supplements

---

- **Absolutely NO smoking, vaping, nicotine patches, nicotine gum, any nicotine products, or secondhand smoke exposure for at least 4 weeks before surgery and 4 weeks after surgery.** Nicotine constricts blood vessels and can cause skin loss, poor healing, and worse scarring.
- **AVOID ALCOHOL for at least 2 weeks before surgery and 2 weeks after surgery.** Alcohol use significantly increases the risk of bleeding, bruising, and swelling.

## Hair

---

- Be very gentle with your hair. Do not pull a comb or brush through the incisions. Doing so may disrupt or open the incision lines. Let water and shampoo rinse crusts loose in the shower.
- Do not color, highlight, perm, or chemically treat your hair for 4 weeks after surgery.

## Questions or Concerns?

---

Please do not hesitate to contact our office at **949.650.8882** (Newport Beach) or **310.467.2180** (Beverly Hills) during business hours. For life-threatening symptoms, always call 911 or go to the nearest emergency room first.

*Healing is a gradual process. We are here to guide you every step of the way.*